



## Isolation, Partial Purification and Characterization of Lipase Enzyme from *Pseudomonas aeruginosa* from Soil and Sewage Waste

P. Srinivasa Chary<sup>1</sup> and Y. Prameela Devi<sup>2\*</sup>

<sup>1</sup>Centre for Biotechnology, Jawaharlal Nehru Technological University Hyderabad (JNTUH),  
Hyderabad, India.

<sup>2</sup>Department of Zoology, Kaktiya University, Warangal, India.

### Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/AJOB/2018/38394

#### Editor(s):

- (1) Paola Angelini, Department of Chemistry, Biology and Biotechnology, University of Perugia, Perugia, Italy.  
(2) Xing Li, Division of Biomedical Statistics and Informatics, Department of Health Sciences Research, Mayo Clinic College of Medicine, USA.

#### Reviewers:

- (1) Carlos Luna, University of Cordoba, Spain.  
(2) Samia Mezaache-Aichour, Université de Sétif, Algeria.  
(3) Meshack Hawi, Jomo Kenyatta University of Agriculture and Technology, Kenya.  
(4) Maria Laura Belaunzaran, University of Buenos Aires, Argentina.  
Complete Peer review History: <http://prh.sdiarticle3.com/review-history/24707>

Received 24<sup>th</sup> November 2017

Accepted 17<sup>th</sup> May 2018

Published 22<sup>nd</sup> May 2018

Original Research Article

### ABSTRACT

A lipase producing *Pseudomonas* sp. is isolated from soil and sewage waste water in Hyderabad. The growth curve of *Pseudomonas aeruginosa* on growth medium is plotted. It is capable of producing the enzyme lipase where the specific enzyme activity is maximum. The optimum hydrolysis conditions for lipase are pH 7.0 to 9.0 and temperature at 45.0 C. These results show that *Pseudomonas aeruginosa* may have potential for industrial sewage waste water treatment and recycling of sewage wastes from the environment.

*Keywords:* Lipase; pH; sewage; temperature and waste water.

\*Corresponding author: E-mail: [prameeladevi@yahoo.co.in](mailto:prameeladevi@yahoo.co.in);

## 1. INTRODUCTION

Fossil fuels are consumed so fast and the planet resources will be going to be exhausted soon. There is a need to develop inexpensive and alternative new technologies to produce sustainable green fuels. Besides hydrolytic reactions lipases undergo esterification reactions or transesterification reactions thereby useful for the production of biodiesel, an alternative energy. The selection of the lipases producing organism that perform transesterification reactions is not easy. There is few biodiesel producing lipases from micro organisms are currently available for industrial purpose [1].

The application of enzymes in the transesterification reaction or process of triglycerides making integration of glycerol to form monoglycerides. There is an increasing yield in the process and also the environmental sustainability of the conventional biodiesel production. This is called as Ecodiesel [2]. Lipases occur widely in nature but microbial lipases are commercially significant. Microbial lipases have wide applications in industries because of their shorter generation time, ease of production (that further enhanced with advancement in fermentation technologies) and ease of manipulation (either genetically or environmentally). Microorganisms that produce lipase are found in diverse habitats such as dairies, vegetable oil processing factories, industrial wastes and soil contaminated with oil. Many microorganisms such as bacteria, yeast and fungi are known to secrete lipases but bacterial lipases are more economical and stable. Several lipases from *Bacillus thermoleoverans*, *B. stearothermophilus*, *B. acidocaldarius*, *thermoacidophilic bacteria* and *Bacillus* sp. strain A 30-1 (alkaliphilic bacteria) have been isolated and characterized. Bacterial lipases are used extensively in food and dairy industry for the hydrolysis of milk, fat, cheese ripening, flavor enhancement and lipolysis of butter fat [3].

Bacterial lipases are extracellular, they are greatly influenced by physico-chemical and nutritional factors, such as temperature, pH, nitrogen & carbon sources, inorganic salts, agitation and dissolved oxygen concentration. The carbon sources accounted for 50% of the production cost in the lipase production. In order to reduce the cost of production, usage of cheaper substrates like oil seed cakes, peanut oil cake, groundnut oil cake, palm oil effluent etc., is

advisable. In further to these substrates, oil mill effluent can also be used as substrate for lipase production since it is rich in oil, suspended solids, organic and inorganic nutrients, carbohydrates etc.

Microbial lipases gained prominence in industry mainly because of its hydrolytic and synthetic activities, high yield, ease of genetic manipulation, regular supply due to absence of seasonal fluctuations and easy cultivation of microbes on inexpensive media [4,5]. Lipases of fungal and bacterial origin are widely used in industrial applications. Some important lipase producing bacterial genera include *Bacillus*, *Pseudomonas* and *Burkholderia*. Lipase/esterase-producing bacteria have been found in diverse habitats such as soil contaminated with oil, dairy waste, industrial wastes, oil seeds and decaying food, compost heaps, coal tips and hot springs [6,7]. The demand for the biocatalysts with novel and specific properties such as specificity, stability, pH, and temperature is increasing (K.E. Jaeger et al.,) [7].

The industrial demand for highly active preparations of lipolytic enzymes continues to stimulate the search for new enzyme sources. Topal et al., [8] in Turkey first isolated lipase from *Trichoderma harzianum* IDM14D but the enzyme was not characterised. It produces some hydrolytic enzymes such as chitinase,  $\beta$ -glucanase, amylase and cellulase. It is also used as a biocontrol agent.

In the present study bacterial species from wastewater was isolated and screened for the best lipolytic strain (for lipase production, oil mill effluent was used as substrate) by optimizing its environmental factors.

## 2. MATERIALS AND METHODS

### 2.1 Screening of Lipolytic Bacteria

Lipase producing microbial cultures was isolated from different soil and sewage industrial sites in Hyderabad. They are enriched by periodic subculturing of samples in Nutrient Broth (NB) media containing 20% (v/v) and 40% (v/v) wastewater in successive samples. The composition of NB medium is (per liter) 5g peptone and 3 g yeast extract. The pH of the medium was adjusted to 7 with 0.1 M NaOH.

## 2.2 Growth Conditions of *Pseudomonas aeruginosa*

The isolation was done by serial dilution of samples on tributyrin agar plates. The composition of tributyrin agar medium per liter is 5 g peptone, 3 g yeast extract, 10 ml tributyrin and 15 g agar. Culture plates were incubated at 30°C. Colonies showing clear zones around them were picked out and purified on tributyrin agar plates, later it has been transferred to agar slants. The organism identified was *Pseudomonas aeruginosa*. Isolates having clearing zone were grown in the liquid culture and the level of lipase production was determined from the cell free culture supernatant fluid. Characterization and identification of the isolate with higher lipolytic activity was carried out biochemically. The identification of the microorganism *P. aeruginosa* was performed using positive oxidase test, the absence of indole production, a positive urease test, the oxidative metabolism of glucose and complete nitrate reduction to nitrogen [9,10] have revealed that the organism is *Pseudomonas aeruginosa*.

## 2.3 Enzyme Production

The composition of production medium used in this study was: (% w/v) peptone -  $\pm 0.2$ ;  $\text{NH}_4\text{H}_2\text{PO}_4$  - 0.1; NaCl - 0.25;  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$  - 0.04;  $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$  -  $\pm 0.04$ ; olive oil - 2.0 (v/v); pH -  $\pm 7.0$ ; 1-2 drops Tween 80 as emulsifier. In 5ml of sterile deionised water, overnight cultures were suspended and used as the inoculum for pre-culture so that an initial cell density to adjust the turbidity of 0.5 McFarland standard is obtained. Submerged microbial cultures were incubated in 500 ml Erlenmeyer flasks containing 100 ml of liquid medium on a rotary shaker (150 rpm) and incubated at 30°C. The culture is incubated for 24 hours and was centrifuged at 10,000 rpm for 20 min at 4°C, the cell free culture supernatant fluid was collected and used as the sources of extracellular enzyme.

## 2.4. Assay of Lipase Activity

Lipase activity was determined spectrophotometrically at 30°C using p-nitrophenol palmitate (pNPP) as substrate. The reaction mixture was composed of 700  $\mu\text{l}$  pNPP solution and 300  $\mu\text{l}$  of lipase solution. The pNPP solution was prepared by adding the solution A (0.001 g pNPP in 1ml isopropanol) to solution B (0.02 g Sodium deoxycholate, 0.01 g gum arabic, 9 ml of 50 mM Tris-HCl buffer, 50  $\mu\text{l}$  Triton X-100, pH 8). The absorbance was measured at

410 nm for the first 2 min of reaction. One unit (1U) was defined as that amount of enzyme that liberated 1  $\mu\text{mol}$  of pNPP per minute under the test conditions. Assay of Lipase was done with p-nitrophenyl fatty acyl esters at various chain lengths. Absorbance reading at 400 nm was taken with the certain time interval. The curve which has a raising behavior was drawn according to absorbance versus minutes data. The highest enzyme activity data assumed 100% and relative enzyme activity estimated for each data. To calculate specific enzyme activity (units/mg protein), Bradford assay was performed. Specific enzyme activity was obtained by dividing enzyme activity value to protein content in milliliter. At optimum pH and temperature one unit releases one nanomole ( $10^{-9}$  mole) of p-nitrophenol per using p-nitrophenyl fatty acyl esters at various chain lengths.

## 2.5 Effect of pH on Lipase Activity

Optimal pH was determined at 30°C in buffer solutions of pH values ranging from 5 to 11 (0.05 M citrate-phosphate, pH 5-7; 0.05 M Tris-HCl, pH 8-9; 0; 0.05 M Glycine - NaOH, pH 11). By the spectrophotometric assay the effect of pH on enzyme stability was analyzed after pre-incubation of 300  $\mu\text{l}$  of enzyme solution for 1 h at 30°C in 700  $\mu\text{l}$  of the above mentioned buffer solutions (pH 5-11). Enzymatic activity was measured according to a standard protocol with pNPP as the substrate.

## 2.6 Effect of Temperature on Lipase Activity

The crude enzyme used for assay was the culture broth after separation of cells and particles. The enzyme was stored at 4°C until used. The optimal temperature for the activity of enzyme was determined by performing assay at different temperatures (30-70°C), at pH 8.0 for 10 min. For determination of temperature stability, the reaction mixtures containing the enzyme in 50 mM Tris-HCl buffer (pH 8.0) was incubated at different temperatures (37, 45, 50, 55, 65 and 70°C) for 3 h and immediately cooled. Residual enzyme activity was measured under standard enzyme test conditions.

## 3. RESULTS AND DISCUSSIONS

### 3.1 Isolation of Pure Cultures of *Pseudomonas aeruginosa*

The isolation was done by serial dilution of samples on tributyrin agar plates. Culture plates were incubated at 30°C. Colonies with clear

zones around them were picked out and purified on tributyrin agar plates, later transferred to agar slants. The organism identified was *Pseudomonas aeruginosa*. *Pseudomonas aeruginosa* is a gram-negative rod shaped bacteria measuring 0.5 to 0.8  $\mu\text{m}$  by 1.5 to 3.0  $\mu\text{m}$ . maximum strains are motile by means of a single polar flagellum. The smooth and mucoid colonies are presumed to play a role in colonization and virulence. Natural isolates from soil or water typically produce a small, rough colony. In general clinical samples yield one or another of two smooth colony types (Fig. 1).



Fig. 1. Culture of *Pseudomonas aeruginosa*

### 3.2 Enzyme Assay of Lipase

One unit (1U) was defined as that amount of enzyme that liberated 1 $\mu\text{mol}$  of pNPP per minute under the test conditions. Assay of Lipase was done with p-nitrophenyl fatty acyl esters at various chain lengths. Absorbance reading at 400nm was taken with the certain time interval. The maximum enzyme activity is obtained.

### 3.3 Effect of pH on Lipase Activity

The effect of pH on the activity of lipase enzyme was determined using four different buffers with pH 3.0 to 12.0. The enzyme was most active at pH 6.0 & 9.0 increasing activity from pH 7-9 can be characterized as an alkalophilic enzyme but high lipase activity at pH 6 shows its activity in acidic conditions. Bacterial lipases are stable between pH 4-11. Lipase obtained in our study was stable at basic conditions from pH 7.0 to 10.0 but its stability was low at acidic pH. The remarkable stability of *Pseudomonas aeruginosa*

lipase in this range has proved it to be a potential alkaline lipase (Fig. 2). The optimal activity of lipases extracted from *Bacillus thermoleovorans* ID1 was at pH 9 [11]. Thermophilic bacteria like *Bacillus* that produce lipases showed optimal activity between pH 7.2–8.5 [11-18]




Fig. 2. Effect of pH on lipase enzyme activity




Fig. 3. Effect of Temperature on lipase enzyme activity

### 3.4 Effect of Temperature on Lipase Activity

The temperature preference of this enzyme reveals higher activity values at temperature from 35 to 45°C. The lower activity at 35°C compared to 45°C, probably is due to the kinetics of enzymatic reaction as the enzyme is more conductive at 45°C. Assessment of the thermostability of lipase was performed by measuring the residual activity at various times, following incubation at different temperatures. *P. aeruginosa* lipases seem to be more

thermostable than others from this genus (Fig. 3). *B. thermoleovorans* ID-1 showed optimal activity between 60–65°C [11,]. *Bacillus sp.* J33 showed optimal lipase activity at 60°C [14], *Bacillus sp.* A30-1 also showed maximum activity at 60°C [18].

#### 4. CONCLUSION

The results obtained in this study show that the isolate *P. aeruginos* has produced a maximum concentration of lipase enzyme. Further studies are needed to enhance lipase production in this strain. Also, *Pseudomonas aeruginosa* lipase is a potential alkaline lipase. The enzyme exhibited maximum activity and stability between pH 7.0 and 10.0 and temperature of 45°C. The maximum enzyme activity is found as 50 units. The remarkable stability of *P. aeruginosa* lipase in this range has proved it to be a potential alkaline lipase similar to other and a candidate for industrial applications such as detergent, leather and fine chemical industries.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Escobar-Niño A, Luna C, Luna D, Marcos AT, Cánovas D, Mellado E. Selection and characterization of biofuel-producing environmental bacteria isolated from vegetable oil-rich wastes. PLoS ONE. 2014;9:e104063. DOI: 10.1371/journal.pone.0104063
- Luna C, Luna D, Bautista FM, Estevez R, Calero J, Posadillo A, Romero AA, Sancho E. Application of enzymatic extracts from a CALB standard strain as biocatalyst within the context of conventional biodiesel production optimization. Molecules. 2017; 22:2025. DOI: 10.3390/molecules22112025
- Arpigny JL, Jaeger KE. Bacterial lipolytic enzymes: 5. classification and properties. Biochem J. 1999;343:177-13.
- Jaeger K, Ransac S, Dijkstra BW, Colson C, Heuvel M, Misset O. Bacterial lipases. FEMS Microbiology Reviews; 1994.
- Hasan F, Shah AA, Hameed A. Industrial applications of microbial lipases. Enzyme Microb. Technol. 2006;39:235-251.
- Vakhlu J, Kour A. Yeast lipases: Enzyme purification, biochemical properties and gene cloning. Elect J Biotechnol. 2006;9: 69-85.
- Wang Y, Srivastava KC, Shen GJ, Wang HY. Thermostable alkaline lipase from a newly isolated thermophilic *Bacillus* strain, A30-1(ATCC 53841). J ferment Bioeng. 1995;79:433-8.
- Topal S, Pembeci C, Borcaklı M. Türkiye'nin tarımsal mikoflorasının endüstriyel öneme sahip bazı enzimatik aktivitelerinin incelenmesi-I: Amilaz, proteaz, lipaz. Turk J Biol. 2000; 24:79-93,.
- Szita G, Biró G. A synthetic culture medium for *Pseudomonas aeruginosa*. Acta Vet Hung. 1990;38:187–194.
- Szita G, Biró G, Tóth G, Pintér V, Fábrián A. Detection of *Pseudomonas aeruginosa* using a synthetic medium (in Hungarian, with English abstract). Magy Allatorv Lapja (Hung Vet J ).1995;50:287–289.
- Lee DW, Kim HK, Lee KW, Kim BC, Choe AC, Lee HS. Purification and characterization of two distinct thermostable lipases from the gram-positive thermophilic bacterium *Bacillus thermoleovorans* ID-1, Enzyme Microb Tech. 2001;29:363–71.
- Schmidt DC, Sztajer H, Stocklein W, Menge U Schmid RD. Screening, purification and properties of a thermophilic lipase from *Bacillus thermocatenuatus*. Biochimica et Biophysica Acta 1994;1214: 43–53.
- Kim HK, Sung MH, Kim HM, Oh TK. Occurrence of thermostable lipase in thermophilic *Bacillus sp.* Strain 398, Biosci Biotech Bioch. 1994;58:961–2.
- Nawani N, Dosanjh NS, Kaur J. A novel thermostable lipase from a thermophilic *Bacillus sp.*: Characterization and esterification studies, Biotechnol Lett. 1998;20:997–1000.
- Lee DW, Koh YS, Kim KJ, Kim BC, Choi, HJ, Kim DS. Isolation and characterization of a thermophilic lipase from *Bacillus thermoleovorans* ID-1. FEMS Microbiol Lett. 1999;179:393–400.
- Dharmsthiti S, Luchai S. Production purification and characterization of thermophilic lipase from *Bacillus sp.* THL027. FEMS Microbiol Lett. 1999;179: 241–6.

17. Imamura S, Kitaura S. Purification and characterization of a monoacylglycerol lipase from the moderately thermophilic *Bacillus* sp. H-257, J Biochem. 2000;127: 419–25.
18. Handelsman T, Shoham Y. Production and characterization of an extracellular thermostable lipase from a thermophilic *Bacillus* sp., J Gen Appl Microbiol. 1994; 40:435–43.

---

© 2018 Chary and Devi; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<http://prh.sdiarticle3.com/review-history/24707>