



Incidence of Root-knot Nematodes in Some Selected Vegetable Crops in Nukkai Irrigation Field Jalingo, Taraba State, Nigeria

D. L. David¹, B. W. Barau¹, J. A. Wahedi^{2*} and Y. N. Ahmed³

¹Department of Biological Sciences, Taraba State University, Jalingo, Nigeria.

²Department of Zoology, Adamawa State University Mubi, Nigeria.

³Department of Agricultural Technology, Federal Polytechnic Bali, Taraba State, Nigeria.

Authors' contributions

This work was carried out in collaboration between all authors. Author DLD designed and supervised the study, wrote the protocol and the first draft of the manuscript. Authors BWB and YNA performed the field study, author JAW managed the analyses of the study. Author BWB managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJOB/2017/33900

Editor(s):

(1) Farjana Sultana, College of Agricultural Sciences, International University of Business Agriculture and Technology (IUBAT University), Bangladesh.

Reviewers:

(1) Abdullahi M. Nuhu, Kaduna Polytechnic, Nigeria.

(2) Mahmoud M. A. Youssef, National Research Centre, Egypt.

Complete Peer review History: <http://prh.sdiarticle3.com/review-history/20173>

Original Research Article

Received 3rd May 2017
Accepted 6th June 2017
Published 24th July 2017

ABSTRACT

A study to determine the incidence of root-knot nematode (*Melodogyne* spp) in some selected vegetables was conducted in Nukkai Irrigation Field of Jalingo, Nigeria. Samples of Okra (*Hibiscus esculentus*), Spinach (*Amaranthus* spp) and Sorrel (*Hibiscus sabdariffa*) were collected at 2-weeks old and at maturity (flowering stage). Their roots were cut off and nematodes (*Melodogyne* spp) were extracted using the Baermann method and identified using the female perineal pattern manual. The results showed that *Melodogyne* spp are incident in the study area as 248 (34.44%) stands out of the 720 stands studied were found to be infested by root-knot nematode. The results also showed that two *Melodogyne* spp, *M. javanica* and *M. incognita* were discovered, and had a total number of 535 individuals. *M. javanica* (345) were predominantly higher than *M. incognita* (190). In all stages of sampling, *Hibiscus esculentus* proved to be more susceptible to infestation, followed by *Amaranthus* spp while the least infested was *Hibiscus sabdariffa* and *M. javanica* appeared to be more abundant than *M. incognita* in the study area.

*Corresponding author: E-mail: wahedijasini@gmail.com, wajasini@yahoo.com;

Keywords: *Amaranthus*; *hisbiscus*; *jalingo*; *melodogyne*; *Nigeria*, *Nukkai*.

1. INTRODUCTION

Nematodes are soil dwelling organisms that constitute one of the largest animal phyla in the world, with over half a million species known. It has been estimated that four out of every five living animals on this planet are nematodes [1]. Nematodes can live as obligate parasites of plants and animals and can also alternate a parasitic life, with a free living life style or can be strictly free [2]. Despite the diversity in their life style and habitat, all nematodes are morphologically, anatomically and developmentally similar [3].

Root-knot nematodes (*Meloidogyne* spp) are minute worm-like animals that are very common in the soil and can invade roots of plants. They have a wide host range, and cause problems in many annual and perennial crops; they occur throughout the world infecting all major crops and causing substantial reduction in yield and quality [4]. There are four important species of the genus; *Meloidogyne arenaria*, *Meloidogyne hapla*, *Meloidogyne incognita* and *Meloidogyne javanica* which are considered as the most economically important species responsible for causing more damage to agricultural produce [5,6].

Vegetables are much more important commodities needed in high demand due to their nutritive value for balanced nutrition in many areas of the world but because of root-knot nematode (*Meloidogyne* spp), it is increasingly very difficult and sometimes impossible to grow important vegetable in the tropics and semi tropical countries [2]. In Africa, the yield of vegetables is relatively lower, for which there are many constrains including prevalence of disease caused by different pathogens [7]. Among various pathogens responsible for the low yield, root-knot nematodes are of considerable economic importance [8] and can cause an annual loss of 22% in the tropics [9]. In addition, these parasites also interact with other disease causing organisms to produce a disease complex [10]; break down resistance against other pathogens and reduce plant tolerance to environmental stress [11,12].

It is against this background that this study was designed to determine the incidence of root-knot nematodes of some vegetables cultivated in Nukkai Irrigation Fields of Jalingo, Taraba State.

2. MATERIALS AND METHODS

2.1 Study Area

Nukkai Irrigation Field is an irrigation farming field in Jalingo, Taraba State, which has an area of about 20 hectares. Jalingo is located at latitudes 8°47' to 9° 01'N and longitude 11° 09' to 11°30'E. Nukkai Irrigation Field is located immediately after the Nukkai Bridge south of the Jalingo-Wukari main road in the city of Jalingo (Fig. 1).

2.2 Sample Collection

All samples were collected using the method adopted by Anwar and McKenry [13]. Samples of Okra (*Hibiscus esculentus*), Spinach (*Amaranthus* spp) and Sorrel (*Hibiscus sabdariffa*) were randomly collected at different growth stages, i.e. seedling stage and matured stage. For each vegetable species, seedling was collected at random when they were aged 2-weeks old and the matured vegetables were also collected immediately they started flowering. All samples collected were washed and cleaned of soil debris. The roots were cut off and kept in the fridge in a plastic container.

2.3 Extraction of Nematodes

Nematodes were extracted from roots of plants using the Baermann funnel extraction method as described by Baermann [14]. In the set-up, a funnel was placed in a stand and filled with water until it reached up to 1cm below the rim. A rubber tube was fixed at the mouth of the funnel whose end was tightened using a clip. It was ensured that formation of air bubbles was avoided. The clip ensured it was well closed and the rubber tube did not leak. The sieves with the sample were hung in the funnel so that the sample was totally submerged, without touching the bottom of the funnel. Nematodes crawled out of the sieve into the water and settled. After a period of 16-72 hours, the nematodes suspension was trapped by opening the squeezer clip, regularly tapping and adding water to increase nematodes vitality.

2.4 Identification and Counting of Nematodes

Nematodes were identified using the female perineal pattern method manual or guide by [15]. During identification, nematodes were counted directly under microscope using the counting dish.




Fig. 1. The map of Jalingo showing the study area (Nukkai irrigation field)

2.5 Statistical Analysis

The collected data was analyzed using descriptive statistics of means, percentages and chi-square.

3. RESULTS

3.1 Incidence of Infestation

The incidence of infestation of plant by root knot nematodes is shown in Table 1. The result revealed that plant parasitic nematodes are incident in the field of study as they are found in all the studied vegetable. Out of the 120 stands

from the three (3) plots sampled, Okra (*Hibiscus esculentus*) had the highest infestation of 60 (50.0%), followed by Spinach (*Amaranthus* spp) which had a total infestation of 52 with an incidence of 43.3% while the least infestation (30) was recorded in Sorrel (*Hibiscus sabdariffa*) with a total incidence of 25.0%. Chi-square test also revealed that there was a significant difference ($p > 0.05$) between the total number of infested and uninfested stands studied (Table 1). Across all plots, plot A showed highest infestation, with an incidence rate of 42.5%, followed by plot B (40.0%), while the least infestation was observed in plot C (35.8%).

The incidence of infestation of matured stage of the vegetable plants by root knot nematodes is presented in Table 2. Matured Spinach (*Amaranthus* spp) had the highest total infestation of 54 stands out of the 120 stands from the three (3) plots sampled with an incidence rate of 45.0%, followed by Okra (*Hibiscus esculentus*) which had a total infestation of 51 with an incidence of 42.5% while the least infestation of 38 was recorded in Sorrel (*Hibiscus sabdariffa*) with a total incidence of 31.7%. Chi-square test showed that there is a significant difference ($p > 0.05$) between the total number of infested and uninfested stands of all the vegetable plants (Table 2). The results also revealed that plot A had the highest infestation rate (40.0%) followed by plot B (39.2%) while the least infestation was recorded in plot C (35.8%), as shown in Table 2.

Table 3, showed the abundance of *Meloidogyne* spp extracted throughout the study. Out of the 720 stands of vegetables studied across the three plots, 285 stands were found to be positively infested. A total of 535 *Meloidogyne* spp of nematode were extracted, out of which *M. incognita* recorded a total of 190 individuals across the three plots with the highest (78 individuals) in Plot B, while the least in Plot C (51). *M. javanica* proved to be more abundant in the study area which had a total of 345 individuals across the three Plots, with Plot A having the highest (115 individuals), followed by Plot C (127 individuals), and the least (103 individuals) was recorded in Plot B (Table 3). The chi-square analysis also revealed that there was no significant difference ($p > 0.05$) between the total number of nematodes extracted across the three plots (Plots A, B and C) but there was a significant difference ($p > 0.05$) between the total number of *M. incognita* and *M. javanica* extracted.

4. DISCUSSION

Vascular feeders have become sedentary endoparasites [16]. They usually damage their hosts by redirecting large amount of energy and nutrients from normal cellular activities into their special feeding sites and developmental activities [16]. They also alter tissues and then disrupt the vascular system, which leads to prevention of water transportation and dissolved nutrients upward to the shoots by the xylem and also the translocation of photosynthesized materials to other parts of the plants by the phloem [17].

Roots severely galled by these nematodes could predispose the plants to root rots which could subsequently lead to shorter life span of the plants [18]. Such galled tissues could become succulent, poorly protected from invasion by other disease pathogens and nutrient rich food substance that can easily be colonized by fungi [18]. Anwar and Mckenry [19] reported that damages by root-knot nematodes could lead to poor growth, decline in quality and yield, and reduced resistance to stresses like drought and disease.

As revealed in this study, it is important to note that, there is an increase in number of infestation in each vegetable crop as it ages. This probably signifies that as the plants aged and increases in size, more spaces are created for more nematodes to infest and colonizes the vegetable crops. In all stages of sampling, *Hibiscus esculentus* proved to be more susceptible to infestation, followed by *Amarantus* spp while the least infested was *Hibiscus sabdariffa*. This probably might be due to selection of host by the parasites which favored the attack or infestation of *Hibiscus esculentus* than the rest of the vegetables. It could also be that, Sorrel (*H. sabdariffa*) had the least number of infested stands probably because of its sour nature which hinders the nematodes from infesting it more.

In terms of abundance, this study identified two species of root-knot nematodes from the study area, *Meloidogyne incognita* and *Meloidogyne javanica*. *M. javanica* appeared to be more abundant than *M. incognita* and this could probably be as a result of the nature of the soil in the study area, which might be more suitable for the survival of *M. javanica* than *M. incognita*. It could also be that *M. javanica* is more resistant to stress and other environmental challenges than *M. incognita* which would generally give *M. javanica* reproductive advantage over its cousin (*M. incognita*). This study is in conformity with the works of Haroon and Zylstru [3], Nagesh et al., [20] and Kayami et al., [9] who all reported that root-knot nematodes, *M. javanica* is abundantly available than the other species. The result of this study also agrees with the work of many researchers among which are Khan [21], Verdejo-Lucas et al. [22], Anwar and McKenry [13] and Kayami et al. [9] who all reported availability or incidence and abundance of root-knot nematode in different regions of the world.

Table 1. Incidence of plant parasitic nematodes at seedling stage in Nukkai irrigation Field, Jalingo (2016)

Plant species	Plot A		Plot B		Plot C		Total	
	+ve	-ve	+ve	-ve	+ve	-ve	+ve	-ve
<i>Amaranthus</i> spp	19(47.5%)	21(52.5%)	13(32.5%)	27(67.5%)	20(50.0%)	20(50.0%)	52(43.3%)	68(55.7%)
<i>Hibiscus esculentus</i>	20(50.0%)	20(50.0%)	25(62.5%)	15(37.5%)	15(37.5%)	25(62.5%)	60(50.0%)	60(50.0%)
<i>Hibiscus sabdariffa</i>	12(30.0%)	28(70.0%)	10(25.0%)	30(75.0%)	08(20.0%)	32(80.0%)	30(25.0%)	90(75.0%)
Total	51(42.5%)	69(57.5%)	48(40.0%)	72(60.0%)	43(35.8%)	87(64.2%)	142(39.4%)*	218(60.6%)*

Note: Values with asterix are statistically significant ($\chi^2_{=0.05}$)

Table 2. Incidence of plant parasitic nematodes at matured stage in Nukkai irrigation Field, Jalingo (2016)

Plant species	Plot A		Plot B		Plot C		Total	
	+ve	-ve	+ve	-ve	+ve	-ve	+ve	-ve
<i>Amaranthus</i> spp	20 (50.0%)	20 (50.0%)	19 (47.5%)	21 (52.5%)	15 (37.5%)	25 (62.5%)	54 (45.0%)	66 (55.0%)
<i>Hibiscus esculentus</i>	18 (45.0%)	22 (55.0%)	16 (40.0%)	24 (60.0%)	17 (42.5%)	23 (57.5%)	51 (42.5%)	69 (57.5%)
<i>Hibiscus sabdariffa</i>	10 (25.0%)	30 (75.0%)	12 (30.0%)	28 (70.0%)	16 (40.0%)	24 (60.0%)	38 (31.7%)	82 (68.3%)
Total	48 (40.0%)	72 (60.0%)	47 (39.2%)	73 (68.8%)	48 (40.0%)	72 (60.0%)	143 (39.7%)	299 (60.3%)

Note: Values with asterix are statistically significant ($\chi^2_{=0.05}$)

Table 3. Abundance of *Melodogyne* spp collected during the studies (2016)

Nematode spp	Plot A	Plot B	Plot C	Total
<i>Melodogyne incognita</i>	61	78	51	190*
<i>Melodogyne javanica</i>	115	103	127	345*
Total	176	181	178	535

Note: Values with asterix are statistically significant ($\chi^2_{=0.05}$)

5. CONCLUSION

In conclusion, the study revealed that at all stages of sampling, *H. esculentus* proved to be more susceptible to infestation, followed by *Amarantus* spp while the least infested was *H. sabdariffa*. Meanwhile, *M. javanica* appeared to be more abundant than *M. incognita* in the study area.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Platt HM. The Phylogenetic systematics of free living nematode. The Ray Society, London. 1994;5.
- Babatola OO. A complete characterization of the four most common species of root-knot nematode. *Meloidogyne* with pictorial key. IM Publication Raleigh, N.C.U.S.A. 2003;20-22.
- Haroon SA, Zylstru C. Rapid identification of genetic relationship of *Meloidogyne incognita* population by polymerase chain reaction RAP markers. Egyptian Journal of Agronomatology. 2003;32(4): 18-20.
- Sasser JN. Economic Importance of *Meloidogyne* in tropical countries. In: Lamberti, F. and Taylor C.E. (Eds). Root-knot nematodes (*Meloidogyne* spp): Systematics biology and control. Academic Press, New York. 1987;359–374.
- Wilson JA. The incidence of root-knot nematodes *Meloidogyne arenaria*, *Meloidogyne incognita* and *Meloidogyne javanica* on vegetables. Plant Disease Journal. 2012;22:12–18.
- Babatola OO, Ogurnunmare OB. Interaction between *Meloidogyne incognita* and *Fusarium oxysporum* on selected bean genotypes. Journal of Nematology. 2012;45(2):20-24.
- Ahmed R. Root-knot of potato in West Pakistan. Plant Disease Reporter. 2012; 57:505-510.
- Hassan SS, Nagesh M. Incidence of root-knot nematode, *Meloidogyne incognita* and *Meloidogyne hapla* in Slovenia. Acta Agriculturae Slovenica. 2011;451–474.
- Kayami MZ, Mukhtar T, Hussain MA. Association of root-knot nematodes (*Meloidogyne* spp) with cucumber in the pothwar region of Punjab Province of Pakistan. International Journal of Biotechnology. 2012;9(1-2):23–29.
- Begun GP. Geographical distribution and infestation of plant parasitic nematodes *Fusarium oxysporum* on selected Bean genotypes. Journal of Nematology. 2012; 45(1):231-233.
- Wheeler TA, Hake KD, Dever JK. Survey of *Meloidogyne incognita* and *Thielavipsis basicola*: Their impact on cotton fruiting and producers management choices in infested fields. Journal of Nematology. 2000;32(4):576–583.
- Barbosa DH, Viera SG, Souza RM, Silva CP. Survey of root-knot nematode *Meloidogyne* spp. in Coffee plantations in the state Rio de Janeiro, Brazil. Nematologia Brasileira. 2004;28(1):43–47.
- Anwar SA, McKenry MV. Incidence and population density of plant-parasitic nematodes infecting vegetable crops and associated yield loss in Punjab, Pakistan. Pakistan Journal of Zoology. 2012;44(2): 327–333.
- Baermann G. Baermann method of root-knot nematode extraction. Methods and techniques for nematology, North Carolina State University and state Agency for International Development, Raleigh. 1948; 18-19.
- Taylor DP, Netscher C. An Improved technique for preparing perineal patterns of *Meloidogyne* spp. Nematologica. 1974; 20(2):107–270.
- Anwar SA. Influence of *Meloidogyne incognita*, *Partrichodorus minor* and *Pratylenca scribneri* in root-shoot growth and carbohydrate partitioning in tomato.

- Pakistan Journal of Zoology. 1995;27: 105–113.
17. Hajera H, Feroza N and Shahina F. Effect of verm and nematode interaction on some biochemical parameters of sunflower. Pakistan Journal of Nematology. 2009;27: 193-201.
 18. Abawi GS, Chen J. Concomitant pathogen and pest interactions. In: Plant and Nematode Interactions (eds. Barker KR, Pederson GA, and Windham GI.), Agronomy Monograph 36. American Society of Agronomy, Madiston, WI. 1988;135–158.
 19. Anwar SA, McKenry MV. Incidence and reproduction of *Meloidogyne incognita* on vegetable crop genotypes. Pakistan Journal of Zoology. 2010;42:135-141.
 20. Nagesh M, Hussaini SS, Chidanandaswamy BS. Incidence of root-knot nematode, *Meloidogyne incognita* on gherkin (*Cucumis sativus*) and yield losses. Indian Journal of Plant Production. 2005;33:309-311.
 21. Khan RM. Distribution of *Radopholus similis* in India, its spread in new regions and analysis of the Nematofauna of Banana Crop Pathosystem. Nematologia Mediterranea. 1999;27(2): 239-246.
 22. Verdejo-Lucas S, Ornat C, Sorribas FJ, Stcheigel A. Species of root-knot nematodes and fungal eggs parasites recovered from vegetables in Almeria and Barcelona, Spain. Journal of Nematology. 2002;34(4):405–411.

© 2017 David et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://prh.sdiarticle3.com/review-history/20173>